

Non-Auditory Effects of Environmental Noise

by Marshall Chasin, AuD. for the Canadian Academy of Audiology

Preamble:

This paper is an overview of the findings in the literature for the subject area of the non-auditory effects due to lower-level environmental noise. Regular reviews have appeared in the literature since the early 1990s (Abel, 1990; Fay, 1991), with the most recent one being van Kamp et al. (2020). Over the past decade, the World Health Organization (WHO) has sponsored a number of excellent reviews on the effects pertaining to stress and cardio vascular effects (Van Kempen, Casas, Pershagen, and Foraster, 2017, 2018; Munzel et al., 2018), sleep disruption patterns (Basner, and McGuire, 2018), cognition (Clark, and Paunovic, 2018), adverse birth outcomes issues (Nieuwenhuijsen, Ristovska, and Dadvand, 2017), and general annoyance (Guski, Schreckenberger, and Schuemer, 2017), and have come out with a number of general recommendations that communities and jurisdictions should consider in their environmental and building planning processes (WHO, 2018).

This paper will serve to present a summary of the relevant references and their findings, while attempting to delineate many of the inherent difficulties in performing such research. This in turn, may provide insight into conclusions stemming from the WHO (2018) general recommendations paper.

Indeed, in a recent paper, where only high quality and controlled research was reviewed, it was argued that the WHO (2018) general recommendations paper, may need to be revisited and updated due to the possibility of confounding variables in previously reviewed studies, as well as the inclusion of other environmental noise sources such as wind turbine noise and rail traffic noise (van Kamp, Simon, Notley, Baliatsas, and van Kempen, 2020).

Introduction:

While it is well-established that prolonged exposure to sound levels at 85 dBA and higher can cause a measurable permanent sensori-neural hearing loss (ISO-1999, 1990; Roberts and Neitzel, 2019) there are few reliable measures of systemic dysfunction for levels below 80 dBA. (There are data showing that there may be a slight measurable hearing loss for a sufficiently long exposure of sound levels between 80 dBA and 84 dBA). Reports of non-auditory effects of noise exposure are abundant in the literature with ramifications that can increase stress, disrupt sleep, and can be implicated as a factor in language and reading development for elementary age students in schools.

Some of the literature is contradictory, some reports demonstrate slight issues, and yet other reports fail to find any systemic effects. Part of the problem is the lack of reliable measures, or measures that have only a loose correlation with a systemic pathology. For example, serum cholesterol measures may show a slight elevation in the presence of long-term noise exposure but this should not be taken as an indicator of eventual cardiac pathology. Another example is

that environmental noise that creates the greatest disruption of sleep may have the lowest annoyance when compared to other noise stimuli (see for example, Elmenhorst et al., 2019).

Only publications that have gone through a rigorous peer academic review will be included. However, despite the research appearing in a peer reviewed publication, there can still be serious methodological problems that call the results, or lack thereof, into question. Many historical studies have been summarized in a 1991 book *Noise & Health* (Thomas H. Fay, Ed., 1991) and also a review article by Abel (1990). Davies and van Kamp (2012) also performed an enlightening review, and the results were similar to earlier publications. For an historical review these references should be consulted. Following is a summary of the more recent research which is delineated under four general headings: “Annoyance”, “Stress and cardiovascular effects of noise”, “Sleep disruptions due to noise”, and “Cognitive/educational issues due to noise”.

An inherent problem with examining the relationship between the effects of low levels (below 85 dBA) of environmental noise and non-auditory sequelae is the line between academic integrity and advocacy may be crossed. While it is true that environmental noise exposure may cause some issues with some people, it cannot necessarily be concluded that this would have long-term detrimental consequences. There are no “smoking gun” reports in the literature that environmental noise levels below 85 dBA will cause permanent issues, and this is true of sleep disruption, increased stress leading to cardiac involvement, or long-term educational/language problems. It may be true that some people are more susceptible to environmental noise, and it may be true that the exposure is “dose-dependent” whereby higher level and/or longer exposure to environmental noise may result in greater annoyance that may have long-term effects, but eventual systemic problems such as cardiac issues may also be related to a host of other factors. Low level environmental noise is just one of many factors that can contribute to annoyance and possible long-term health issues.

This review will not concern itself with non-human reactions to environmental noise such as that concerning marine life and other wildlife (see for example, Guan, Scholik-Schlomer, and Pearson-Meyer, 2018 for a review). Many acoustical associations such as the Acoustical Society of America and the Canadian Acoustical Association have had long-standing committees to investigate the effect of marine and environmental noise on migration patterns, reproductive patterns, and even vocalization changes in animals.

General Overview:

A: Annoyance:

Annoyance is primarily a subjective attribute of a sound or vibration. Annoyance can be related to the uniqueness of a sound, its physical spectral and temporal characteristics, and even whether a person has control over the sound. (Bronzaft and Madell, 1991). Miller (1974) stated that noise “has its base in the unpleasant nature of some sounds, in the activities that are disturbed or disrupted by noise, in the physiological reactions to noise, and in the response to the meaning or messages carried by the noise.”

Traditionally “annoyance” has resisted a formal definition since it appears to be the result of many intrinsic and extrinsic factors, can be dose dependent, and can be highly variable among different groups and ages.

Nevertheless, annoyance can be related to some objective measures. This was the basis of much research in the 1950s, 1960s, and 1970s, where the unit “noys” was suggested as a possible measure of annoyance. Spectrally, annoyance appears to be similar to a dB A-weighted noise except that higher-frequency sounds above 1000 Hz contribute more to a sense of annoyance than lower frequencies and can be shown to correlate with the D-weighted filter on sound level meters (Kryter, 1970). Despite this pioneering work, annoyance (and its various measures) remains a highly variable phenomenon with sometimes poor correlations, such as where noise that creates the greatest disruption in sleep, may have a low level of annoyance (see for example, Elmenhorst et al., 2019).

Fields (1993) examined a number of personal and situational variables that could potentially be an issue when trying to predict the noise annoyance in residential settings. “The balance of the evidence from 464 findings drawn from 136 surveys suggests that annoyance is not affected to an important extent by ambient noise levels, the amount of time residents are at home, the type of interviewing method, or any of the nine demographic variables (age, sex, social status, income, education, home ownership, type of dwelling, length of residence, or receipt of benefits from the noise source.” (p. 2753). However, he found that annoyance is related to the amount of isolation from sound at home and to five attitudes - fear of danger from the noise source, noise prevention beliefs, general noise sensitivity, beliefs about the importance of the noise source, and annoyance with non-noise impacts of the noise source. This author also noted that there was a small percentage of people who were very annoyed even by low environmental noise levels.

Miedemaa and Voss (1999) examined a number of previous studies where the number of respondents ranged from 15,000 to 42,000 people. They examined the demographic variables including sex, age, education level, occupational status, size of household, homeownership, dependency on the noise source, and use of the noise source. They also looked at two attitudinal variables- noise sensitivity and fear of the noise source. It was found that fear and noise sensitivity have a large impact on annoyance but that demographic factors are much less important. Noise annoyance is not significantly related to gender, but is related to age. The effects of the other demographic factors on noise annoyance are “very small.”

Van Gerven et al. (2009), in examining the effects of self-reported annoyance from environmental noise across a lifespan, found that of almost 63,000 people between the ages of 15 and 102, there was an age-dependent response. The results revealed an inverted “U” pattern with the youngest and the oldest respondents reporting the lowest levels of annoyance and those around 45 years of age, reporting the greatest annoyance. The authors suggested that a linear relationship over the age span would not be useful in modelling annoyance and could explain the variable results of some earlier studies.

In studying the effects of longitudinal effects of changes in road traffic noise, using questionnaires, measures of annoyance, activity disturbances, and psycho-social well-being were

evaluated by Ohrstrom (2004). Adverse effects of long-term exposure to a high volume of road traffic were studied in socio-acoustic surveys in over a two-year period where traffic had been decreased by 90% from 25,000 cars/day to about 2400 cars/day. This resulted not only in a large decrease in annoyance and activity disturbances, but also in a better general well-being. “To be able to use the outdoor environment and to have the possibility to keep windows open is essential for general well-being and daily behavior, which implies that access both to quiet indoor and outdoor sections of the residency is of importance for achievement of a healthy sound environment.” (p. 719). As a fitting last entry in this section, Ohrstrom went on to say that “future studies should focus more on ‘softer’ health outcomes and well-being ... and preferably be performed in connection with traffic abatement measures”.

B: Cardiovascular effects of noise:

This title is not only apropos for this section but also the name of a chapter by Lawrence Raymond in *Noise & Health* edited by Thomas H. Fay for the New York Academy of Medicine (1991). In summarizing some of the limited generalizations that potentially could be made, he stated (p. 28):

- “1. Abnormal blood pressure levels can be induced by high noise levels in some animal species, and in humans under some laboratory conditions. The effects in animals have been long-lasting in some studies.
2. The relevance of these finding to human subjects is unclear....it is not known whether controlling noise to levels which do not usually cause hearing loss ... leaves any residual risk of causing hypertension...
3. ... the intensity-duration relationship has not been studied.”

Richard Sloan (1991), in summarizing the potential cardiovascular effects due to noise exposure, went on to write that despite many inconsistencies in the literature, the exposure to noise suggests that (1) exposure to noise is associated with elevations in blood pressure, primarily diastolic blood pressure, and (2) that in general, this effect appears not to habituate over time”. (p. 23). Sloan related these findings to the potential explanation of noise-induced vasoconstriction and the effects on the central blood system.

Some of the methodological and research issues that plagued earlier research (as far back as 1946) have still not been adequately resolved. High levels of subject variability, individual susceptibilities with both psychological and possibly genetic differences, and lack of physical objective measures that possess a high correlation with long-term cardio-vascular sequelae, remain problematic. Following is a brief overview of some of the approaches, but perhaps only the molecular analysis study of Munzel et al. (2018) shows any real promise of demonstrating any relationship between low level environmental noise and potential cardiovascular effects.

Basner et al. (2014) attempted to describe, in a review paper, the physiological bases for the effects of lower levels of noise on the body. They describe a generalized stress model whereby

contributions are made by both conscious stress brought about by an emotional response due to perceived discomfort (“indirect pathway”), and non-conscious physiological responses between the central auditory pathway and different portions of the central nervous system (“direct pathway”). In this model long-term chronic exposure can create an allostatic load altering a person’s homeostatic conditions. This in turn can affect the metabolism and the cardiovascular system. (Allostatic load refers to the cumulative burden of chronic stress and life events. When environmental stress exceeds a person’s ability to cope then allostatic load ensues). This “stress response” is well-known in the literature and can be implicated in people who report tinnitus, and generalized fear of loud sounds such as misophonia (e.g., Davis and El Rifaie, 2000)

Walker et al. (2016) sought to examine 10 male subjects on multiple visits over 41 days in a well-controlled laboratory environment. The subjects were exposed to noise levels (below 85 dBA) and the noise was either centred in the 31.5-125 Hz region or the 500-2000 Hz region. Ambulatory ECG, various blood pressure measures, and saliva-based cortisol measures were assessed before and after exposure. Of all of the measures, only Heart Rate Variability (HRV) was found to show a statistically significant change, but only for the lower frequency (31.5-125 Hz) stimulus. There was no statistical evidence for the blood pressure and saliva cortisol measure differences. Although they postulated an allostatic load issue and a generalized stress response, these mechanisms were only hypothetical and no direct data was supplied to support this conclusion.

Michaud, Feder, Keith, et al. (2016) assessed stress reactions associated with wind turbine noise exposure using self-reported and objective measures. Over 1200 participants, between the ages of 18-79 who lived between 0.25 km and 11.22 km from wind turbines had exposures of up to 46 dBA outdoor levels of wind turbine related noise exposure, were assessed. Regression modeling was only able to account for 21-33% of the variance and many of their measures did not achieve statistical significance. This included measures of the Perceived Stress Scale, hair cortisol concentrations, resting blood pressure levels and heart rate. Of these measures, only the subjective Perceived Stress Scale scores were positively, but weakly, related to cortisol concentrations and resting heart rate. Regardless of the measured outdoor noise levels from the wind turbines, diastolic blood pressure appeared to be only slightly higher among participants who were highly annoyed by blinking lights on turbines. “Collectively, the findings do not support an association between exposure to wind turbine noise up to 46 dBA and elevated self-reported, and objectively defined measures of stress.” (p. 1467)

To examine potential changes at the molecular level, specifically looking at Reactive Oxygen Species (ROS), Munzel et al. (2018) demonstrated that aircraft noise exposure during nighttime can “induce endothelial dysfunction in healthy subjects and is even more pronounced in coronary artery disease patients. Importantly, impaired endothelial function was ameliorated by acute oral treatment with the antioxidant vitamin C, suggesting that excessive production of ROS contributes to this phenomenon.” (p. 873). They also describe an animal model of aircraft noise exposure characterizing the underlying molecular mechanisms leading to noise dependent adverse oxidative stress-related effects on the vasculature.

Van Kempen, Casas, Pershagen, and Foraster, M. (2018), in looking at a review of available data concluded that “The results of the current review show that at this moment, not enough studies of good quality are available that investigated the impact of noise on the cardiovascular and metabolic system. The plausibility of an association calls for further efforts with improved research methodology. In order to improve the quality of the existing evidence, more studies with a cohort or case-control design are needed.... We also recommend that more well-designed studies on health effects in relation to exposure to wind turbines and rail traffic noise are set up and carried out.” (pp. 12-13)

C: Sleep disruptions due to noise:

Similar to the studies concerning the interaction between environmental noise and cardiovascular/stress relationships, the study of sleep disruptions due to low-level environmental noise can be highly problematic. Low-level environmental exposure can be viewed as more problematic with some groups of people, and indeed, during some hours of the day with the 6 PM -9 PM time period surprisingly being the most bothersome (Bullen and Hede, 1983). And many of the studies that have been reported in the literature do not touch on the effects of more recent wind turbine noise exposure, which may turn out to be problematic in some conditions and with some population groups.

Jurriens and colleagues (1983) performed the “1000 night joint European field study of noise effects” and found that indeed subjectively the sleep quality was affected negatively after noisy nights, but that the results were highly variable. They did find slight decreases in REM (Rapid Eye Movement) sleep during noisy nights, but only by about 6.5 minutes. There were however no clear implications for long-term systemic implications.

Historically, the study of sleep disruptions were based on subjective reports as well as changes in REM vs. non-REM sleep fractions as well as alterations in the subjects’ EEG activity, and overall EEG power. However caution should be exercised when interpreting “overall” or “complete sleep time” measures since there can be many transient changes that affect the final number. In laboratory settings many studies used white noise that may not be appropriate for generalization where actual environmental noise may be intermittent or restricted to certain frequency regions. To further complicate things, environmental noise that creates the greatest disruption of sleep may have the lowest annoyance when compared to other noise stimuli (Elmenhorst et al., 2019).

It is also not clear from the research whether the data can be extrapolated equally to all groups in society. For example, increased age is correlated with increased arousal from sleep due to low level environmental noise, as are shift workers, especially if alcohol and tobacco are consumed (Pollak, 1991).

In any insightful study, Elmenhorst et al. (2019) examined three environmental noise sources that they considered to significantly contribute to general annoyance: air, road, and railway traffic noise. However, these findings may not be transferable to physiological reactions during

sleep which are considered to decrease nighttime recovery and might mediate long-term negative health effects. They found that measures of annoyance and studies on sleep awakenings were not always correlated. For example, studies on awakenings from sleep indicate that railway noise, while having the least impact on annoyance, may have the most disturbing properties on sleep compared to aircraft noise. The impact of over 100,000 noise events on assessed awakenings were analyzed in 237 subjects. Results showed that the probability to wake up from equal maximum sound pressure levels (SPL) increased in the following order: *aircraft < road = railway noise*. That is, while the probability of waking from road and railway noise exposure was roughly equal, aircraft noise at 70 dB SPL was 7% *less* likely to disrupt sleep as compared with the other two noise sources. They concluded that the three major traffic noise sources differ in their impact on sleep. The order with which their impact increased was opposite to what was compared to that found in annoyance surveys. They state that “It is thus important to choose the correct concept for noise legislation, i.e., physiological sleep metrics in addition to noise annoyance for nighttime noise protection.” (p. 10).

In an older, but nevertheless insightful study, Ando and Hattori (1977) studied the effect of airport noise on unborn fetuses (groups 1, 2, and 3) and in newborns (group 4). The environmental noise levels were higher for this study (95 dBA) and there is some question whether these responses are the result of systemic and metabolic changes from the mother or changes directly to the fetus. The reactions of babies to aircraft noise were studied by means of electroplethysmography (PLG) and electroencephalography (EEG). It was found that the babies whose mothers had moved to the area around the Osaka International Airport before conception (group 1) or during the first five months of pregnancy (group 2) showed little or no reaction on PLG and on EEG to aircraft noise. In contrast, babies whose mothers had moved closed to the airport during the second half of the pregnancy or after birth (groups 3 or 4) and the babies whose mothers lived in a quiet living area (control group), reacted to the same auditory stimuli. The babies in groups 1 and 2 showed different responses depending on whether the auditory stimuli were aircraft noise or music. “Abnormal PLG and EEG were observed in the majority of babies living in an area where noise levels were over 95 dBA. This suggests that the deep sleep of the babies living in such an area was disturbed, but it is not clear whether there would be any long-term effects from these exposures.” (p. 199).

Amundsen, Klæboe, and Aasvang (2013) describes the Norwegian facade insulation study which includes modifications to over 2500 dwellings that serve to reduce the indoor noise levels. In Norway, about 30% of the population lives near main arterial roads with in-door noise levels being on the order of 55 dBA. In an attempt to remediate this situation, a program was implemented in 2004 and 2005 for more than 2500 dwellings with improved windows, building sound-insulation facades, and in some cases, quieter home ventilation systems. The facade-insulating measures reduced indoor noise levels by 7 dB on average. Before the intervention, 43% of the respondents were highly annoyed by noise. Half a year after the intervention, the proportion of respondents who were highly annoyed by road traffic noise had been significantly reduced to 15%. In a follow-up study two years later, they found that the number of self-reported annoyances from the outdoor noise level remained lower, and that the reduction in the respondents’ self-reported sleep disturbances also remained stable from the first to the second

post-intervention study. This suggests that there is some statistical support for this type of assessment. In the control group, there were no statistically significant differences in annoyance between the preintervention and the two post-intervention studies. This study indicated a reduction from 43% to 15% in annoyance level (on a questionnaire) and was related to a 7 dB noise reduction level.

Smith et al. (2017) evaluated the relative contribution and possible interaction effects of vibration and noise from railways on physiological sleep outcomes. Of importance is the finding that both of these environmental sequelae of railways are important. “Vibration from railway freight often accompanies airborne noise, yet this data is almost totally absent in the existing literature.” (p. 3262). In an experimental investigation, 23 participants, each sleeping for six nights in the laboratory, were exposed to 36 simulated railway freight pass-bys per night with vibration alone, noise alone (49.8 dBA), or both vibration and noise simultaneously. A fourth exposure night involved 52 pass-bys with concurrent vibration and noise. Sleep was measured with polysomnography. Cardiac activity was measured with electrocardiography. The probability of awakenings was greater following all exposures, including vibration alone, than spontaneous reaction probability ($p < 0.05$). The effects of vibration exposure and noise exposure on changes of sleep stage and arousals were directly additive. This implies that the effects of vibration and the noise were processed separately in the brain, but the mechanism(s) were not investigated in this study. The findings show that vibration and the noise level, are both important when considering the impact of railway freight on sleep.

Rocha et al. (2019) reported on a survey of sleep disruption for people living near the Atlanta International Airport and items included questions about sleep quality, sleep disturbance by noise, noise annoyance, coping behaviors, and health. Approximately 10% of over 3000 surveys were returned. Some of the responses was that the calculated outdoor nighttime aircraft noise (“Lnight”) was significantly associated with lower sleep quality ($p < 0.05$), trouble falling asleep within 30 min ≥ 1 /week ($p < 0.01$), and trouble sleeping due to awakenings ≥ 1 /week ($p < 0.05$). The Lnight level was also associated with increased prevalence of being highly sleep disturbed ($p < 0.0001$) and highly annoyed ($p < 0.0001$) by aircraft noise. The Lnight level was associated with several coping behaviors such as often or always closing their windows ($p < 0.01$), drinking alcohol ($p < 0.05$), using television ($p < 0.05$), and using music ($p < 0.05$) as sleep aids. There was no significant relationship between the Lnight level and self-reported general health or likelihood of self-reported diagnosis of sleep disorders, heart disease, hypertension or diabetes.

In another study performed by the same group, Bartels et al. (2021), studied the impact of nighttime (Lnight) road traffic noise, bedroom window orientation, and work-related stress on subjective sleep quality on almost 4000 working women. Sleep was assessed using a single question on general sleep quality and four questions on specific sleep problems (poor sleep vs. no poor sleep). Work-related stress was assessed by questions pertaining to job strain and effort-reward imbalance. Night time exposure to road traffic noise was assessed regarding whether the bedroom window faced onto a quiet, medium, or high traffic street noise levels (modelled as <45 dBA, 45-50 dBA, or >50 dBA respectively). Poor sleep was associated with job strain and effort-reward imbalance. The prevalence of poor sleep did not increase with increasing Lnight.

Bedroom windows that faced onto a quiet street had a protective effect on sleep in each Lnight category. They found a non-significant trend for an additive interaction between bedroom window Lnight exposure and job strain, and concluded that noise levels modelled for the most exposed location likely overestimated the actual exposure and thus may not be a precise predictor of poor sleep. Bedroom window Lnight exposures may be a more relevant factor. Although this did not achieve statistical significance the authors queried whether potential additive interaction effects between bedroom window orientation and job strain should be considered when interpreting epidemiological study results on noise-induced sleep disturbances.

D: Cognitive/educational issues due to noise:

Unlike the subject areas of the effects of noise on stress and sleep disruptions, the effects of low-level environmental noise on cognition and educational/language issues are more straightforward. While it is true that some “well-controlled” laboratory-based studies of the 1960s showed little or no effect, others showed adverse effects on some tasks (e.g., Wyon, 1968 as reported by Bronzaft, 1991). But when switching the research paradigm to a more natural “in situ” setting, more educational measures have achieved statistical significance, especially when other factors such as psycho-social aspects are controlled for. It has been argued that the natural settings correlational studies would be better to draw educational and environmental conclusions from rather than (perhaps limited), but well-controlled laboratory studies. (Bronzaft, 1991).

As far back as the early to mid-1970s, studies in a natural environment started to appear. Two in particular can be metaphorically referred to as “smoking gun” studies that show clear results. There is however no evidence that decreased performance on reading or writing assessments will have any long-term effects.

Cohen, Clark, and Singer (1973) were able to show that children who lived in lower floors of apartments, nearer to noisy expressways, performed poorer on school reading tasks than matched children who lived in higher altitude apartments, farther from environmental noise sources.

Bronzaft and McCarthy (1975) published a study with the telling name “The effect of elevated train noise on reading ability”. The school that was used had half of the classrooms facing noisy train tracks and the other side of the school was quieter. The students were matched for socio-economic and other factors. During a typical school day, 80 trains passed the school with levels of 89 dBA and when quiet, the classroom noise levels were 59 dBA. Bronzaft (1991) states that “Students on the noisy side did more poorly on reading achievement tests than did those on the quiet side of the building. The children in the sixth grade who were exposed to the noise were found to be a year behind their counterparts on the quiet side” (p. 88).

Of interest in this last study, noise abatement measures were performed both by the transit authority (rubber vibration isolator pads were installed on the tracks) and the school (acoustic tiling installed in the noisiest classrooms adjacent to the train tracks), resulting in a 6-8 dB reduction in noise levels. In a follow-up study, published 6 years later, after the noise abatement

measures were undertaken, the reading levels were assessed for children on both sides of the school, and found them to be comparable (Bronzaft, 1981).

In the first of four studies by Dockrell and Shield (2004) titled “Children’s perceptions of their acoustic environment at school and at home”, this group assessed the question regarding whether children could be a good judge of the effects of certain environmental noise levels. Over 2000 children completed a questionnaire designed to discriminate different classroom listening conditions, the noise sources heard at home and at school, and their annoyance by these noise sources. Teachers also completed a questionnaire about the classroom noise environment. Indeed, children were able to discriminate between situations with varying amounts and types of noise and their responses accounted for 45% of the variance in the school environment noise sources. External L_{Amax} levels were a significant factor in reported annoyance, whereas external L_{A90} and L_{A99} levels were a significant factor in determining whether or not children hear sound sources. This study demonstrates that children can be sensitive judges of their noise environments and that the impact of different aspects of noise needs to be considered.

In the same year, these same researchers (Shield and Dockrell, 2004) published data on external and internal sound levels in a range of schools in the United Kingdom. Of interest is that they were able to delineate a number of physical changes (such as window glazing) that could ameliorate the noise levels. They examined 142 primary schools that were away from airports or airplane flight paths and found that 86% of these schools had outside environmental levels of 57 dBA. The children reported that they were bothered only while performing quieter activities in the classroom and not while the louder activities were being performed. While they did examine the age of the school and factors such as window glazing, their results were inconclusive.

In the third study of their series in the noise and education analysis of primary schools in the United Kingdom, Shield and Dockrell (2008) examined the effects of classroom noise on the academic attainments of primary school children. “External noise was found to have a significant negative impact upon performance in measures of literacy, mathematics, and science; the effect being greater for the older children.” (p. 133). The analysis suggested that children are particularly affected by the noise of individual external events. Other factors that significantly affected the children’s test scores were internal classroom noise levels, and background levels being significantly related to test results. Even when the data was corrected for social-economic and special needs factors, the negative relationship between performance and noise levels was maintained.

In their fourth and final study in the United Kingdom, this same group studied reading comprehension levels and vocabulary learning tasks in adolescent students (Connolly, Dockrell, Shield, Conetta, Mydlarz, and Cox, 2019). In a study of almost 1000 English high school pupils (564 aged 11 to 13 years and 412 aged 14 to 16 years) reading tasks on laptop computers were completed while the students were exposed to different levels of classroom noise played through headphones. The number of questions attempted, times taken to read the texts and to answer questions were recorded, as well as correct answers to different types of question. Two similar experiments were performed with classroom noise at levels of 50 and 70 dB L_{Aeq}; and the second at levels of 50 and 64 dB L_{Aeq}. The results showed that the performance of all pupils

was significantly negatively affected in the 70 dB LAeq condition, for the number of questions attempted and the accuracy of answers to factual and word learning questions. However, while the answer was clearer for the younger group of children and showed statistical significance, it was more difficult to discern whether the effects at 64 dB LAeq, had a detrimental effect upon the older pupils.

And finally, as a result of the COVID-19 pandemic, where UNICEF estimates that 1.6 billion children across the world have had their education impacted, and have attempted to continue their learning at home, the study by Chere and Kirkham (2021) has some implications for home-based study and environmental noise levels. 129 students, aged 11–18 took part online after passing an online hearing screening test. The students filled out a sociodemographic questionnaire with questions designed to assess their socio-economic and other factors. This was followed by a home environment and noise questionnaire. Three executive function tasks -the Flanker, the Backward Digit Span, and the Wisconsin Card Sorting Test- were completed while listening to either white noise or a home-like environmental noise. Participants were then stratified into quieter and noisier homes for statistical analyses of the results. Measures of the home environment significantly correlated with individual perceptions of the noise and the task performance. Students in this age group, coming from noisier homes were more likely to report that they studied in a noisy room and that they were annoyed by noise when studying. In terms of noise and task performance, the Flanker task revealed that while the older students in the group were more “efficient” than their younger peers, they appeared to lose that advantage if they lived in noisier homes. Reaction times for younger students from noisier homes were less impacted by accuracy compared to their peers from quieter homes, though there was no difference for the older adolescents. “This evidence suggests that higher in-home noise levels lead to higher rates of annoyance and may be hindering home-learning, with both younger and older adolescents being impacted. Furthermore, the long-term effect of in-home noise on adolescent executive function task performance indicates that these findings transcend the pandemic and would influence in-school learning.” (p. 1).

World Health Organization (WHO) Environmental Noise Guidelines for the European Region (2018):

Based on thorough reviews of the available data, the World Health Organization (WHO, van Kempen et al., 2018) has listed some environmental noise limits where non-auditory systemic effects may begin to be observed. The 2018 WHO document was based on a more recent review of meta-studies and reviews from other sources, and many of the authors of the previous reports contributed to the WHO document. All 9 WHO reports are referenced at the end of this document. The 2018 WHO consensus document was based on 7 systematic studies from 2000 to 2014. Van Kamp et al. (2020) reviewed more recent studies and controlled for confounding and methodological issues from the 2018 review. Among the findings were high quality studies linking wind turbine noise to sleep alteration patterns, but that more study needs to be performed for long term effects of environmental noise on cardiovascular dysfunction. Following is a chart, adapted from WHO (2018) summarizing some of the health effects of different average outdoor night noise levels (dB Leq):

< 30 dB Leq	Although individual susceptibilities and circumstances may differ, it appears that up to this level, no substantial biological effects are observed.
30-40 dB Leq	A number of effects on sleep are observed from this range: body movements, awakening, sleep disturbance. The intensity of the effect depends on the nature of the source and the number of the events. Vulnerable groups are more susceptible. A level of 40 dB Leq is equivalent to the lowest observed adverse effect level for night noise.
40-55 dB Leq	Adverse health effects are observed among the exposed population. Many people have to adapt their lives to cope with the noise at night. Vulnerable groups are more severely affected.
>55 dB Leq	This situation is considered increasingly dangerous for public health. Adverse health effects occur frequently, a sizeable proportion of the population is highly annoyed and sleep-disturbed. There is some evidence that the risk of cardiovascular disease increases.

Concluding remarks:

A review of the non-auditory effects of lower-level environmental noise on a number of systems has been reviewed. This is admittedly an extremely difficult area of study with a range of many potentially confounding factors and measures that either have poor test-retest reliability or have a poor correlation with sleep disruption, cognitive and educational issues, stress, and general annoyance. For example, noise that may cause the greatest sleep disruptions may be judged as a noise source that is associated with relatively little annoyance.

There are also a range of other either uncontrolled (or unconsidered) phenomena where some people, in some circumstances, and with some types of noise, are more susceptible to its effects than are others. This may be a dose-related effect, or even a genetic pre-disposition, ... or not-the literature is unclear on this. And to be fair, this same issue of differential “susceptibility” is observed for some section of the population even for high levels of noise and/or music exposure for measurable and permanent hearing loss, where again, the reasons are not well-known.

There are no “smoking gun” studies, or set of studies, that when taken together definitively describe predictable and associated effects for low level noise exposures. While many studies do demonstrate changes in some measures, there is no clear evidence that these changes will have long-term consequences.

While it would be tempting to say the sentence “more research needs to be performed”, given the inherent difficulties in this area of study, it would be unrealistic to conclude that at some point in the near future, definitive and predicted results would be able to be obtained. All that can realistically be said is that for a range of low-level environmental exposures, some people are quite susceptible to its effects, whereas other are not, and that these effects can include sleep disruption, potentially cognitive and educational issues, and an increase in stress and annoyance.

Acknowledgements:

I would like to acknowledge four external readers who provided constructive comments to the manuscript: Dr. Arline Bronzaft, Dr. Bob Harrison, and Dr. Steve Aiken, and Dr. Bill Hodgetts.

References:

- Abel, S.M. (1990). The extra-auditory effects of noise and annoyance: an overview of research. *The Journal of Otolaryngology, Supplement 1*: 1-13.
- Amundsen, A.H., Klæboe, R., and Aasvang, G.M. (2013). Long-term effects of noise reduction measures on noise annoyance and sleep disturbance: The Norwegian facade insulation study
Citation: *The Journal of the Acoustical Society of America* 133, 3921; doi: 10.1121/1.4802824
- Ando, Y., and Hattori, H. (1977). Effects of noise on sleep of babies. *The Journal of the Acoustical Society of America* 62, 199; doi: 10.1121/1.381482.
- Bartels, S., Ögren, M., Kim, J-L., Fredriksson, S., and Persson Waye, K. (2021). The impact of nocturnal road traffic noise, bedroom window orientation, and work-related stress on subjective sleep quality: results of a cross-sectional study among working women. *International Archives of Occupational and Environmental Health*. 94:1523–1536
<https://doi.org/10.1007/s00420-021-01696-w>
- Basner, M., Babisch, W., Davis, A., Brink, M., Clark, C., Janssen, S., and Stansfeld, S. (2014). Auditory and non-auditory effects of noise on health. *The Lancet*, vol. 383, 1325-1332.
- Bronzaft, A.L. “The Effects of Noise on Learning, Cognitive Development, and Social Behavior, in *Noise & Health*. (Thomas H. Fay, Ed.). The New York Academy of Medicine, New York, New York. 1991. (87-92).
- Bronzaft, A.L. (1981). The effects of a noise abatement program on reading ability. *Journal of Environmental Psychology*. 1: 215-222.9
- Bronzaft, A.L., and Madell, J.R. (1991). Community response and attitudes toward noise. in *Noise & Health*. (Thomas H. Fay, Ed.). The New York Academy of Medicine, New York, New York. 1991. (93-99).
- Bronzaft, A.L., and McCarthy, D.P. (1975). The effect of elevated train noise on reading ability. *Environment and Behavior*, 7: 517-528.
- Bullen, R.B., and Hede, A.J. (1983). Time-of-day corrections in measures of aircraft noise exposure. *Journal of the Acoustical Society of America*, 73, 5, 1624-1630.
- Chere, B., and Kirkham, N. (2021). The Negative Impact of Noise on Adolescents’ Executive Function: An Online Study in the Context of Home-Learning During a Pandemic, *Frontiers in Psychology*, 22 September. <https://doi.org/10.3389/fpsyg.2021.715301>

Cohen, S., Glass, D., and Singer, J. (1973). Apartment noise, auditory discrimination and reading ability in children. *Journal of Experimental Social Psychology*, 9: 422-437.

Connolly, D., Dockrell, J., Shield, B., Conetta, R., Mydlarz, C., and Cox, T. (2019). The effects of classroom noise on the reading comprehension of adolescents. *The Journal of the Acoustical Society of America* 145, 372; doi: 10.1121/1.5087126

Davies, H., and van Kamp, I. (2012). Noise and cardiovascular disease: a review of the literature 2008-2011. *Noise and Health*, 14, 287-291.

Davis, A., and El Refaie, A. (2000). The epidemiology of tinnitus, in Tyler, R. (ed.). *Tinnitus Handbook*. Hampshire, Cengage Learning. 1-23.

Dockrell, J.E., and Shield, B. (2004). Children's perceptions of their acoustic environment at school and at home. *The Journal of the Acoustical Society of America* 115, 2964.; doi: 10.1121/1.1652610

Elmenhorst, E-M., Griefahn, B., Rolny, V., and Basner, M. (2019). Comparing the Effects of Road, Railway, and Aircraft Noise on Sleep: Exposure-Response Relationships from Pooled Data of Three Laboratory Studies. *International Journal of Environmental Research and Public Health*, 16, 1073; doi:10.3390/ijerph16061073.

Fields, J.M. (1993). Effect of personal and situational variables on noise annoyance in residential areas. *Journal of the Acoustical Society of America*. 93, 5, 2753-2763.

Guan, S., Scholik-Schlomer, A.R., and Pearson-Meyer, J. (2018). A brief history of our understandings on underwater noise impacts to marine life and the evolution of its regulatory process in the United States. *The Journal of the Acoustical Society of America*, 143, 1767 <https://doi.org/10.1121/1.5035786>.

Jurriens, A.A., Griefahn, B., Kumar, A., Vallet, M., and Wilkinson, R.Y. (1983). An essay in European research collaboration: Common results from the project on traffic noise and sleep in the home. In *Proceedings of the Fourth International Congress on Noise as a Public Health Problem* (Turin), Vol. 2, ed. G. Rossi, Milan: Edizioni Tecniche a cura del Centro Ricerche e Studi Amplifon.

Kryter, K.D. (1970). *The effects of noise on man*. Academic Press, New York.

Miedemaa, H.M.E., and Vos, H. (1999). Demographic and attitudinal factors that modify annoyance from transportation noise. *Journal of the Acoustical Society of America*. 105, 6.

Michaud, D.S., Feder, K., Keith, S.E. et al. (2016). Self-reported and measured stress related responses associated with exposure to wind turbine noise. *The Journal of the Acoustical Society of America* 139, 1467 (2016); doi: 10.1121/1.4942402.

Miller, J.D. (1974). Effects of noise on people. *Journal of the Acoustical Society of America*. 56, 3, 729-764.

Munzel, T., Sørensen, M., Schmidt, F., Schmidt, E., Steven, S., Kroller-Schon, S., and Daiber, A. (2018). The Adverse Effects of Environmental Noise Exposure on Oxidative Stress and Cardiovascular Risk. *Antioxidants & Redox Signaling*. Volume 28, Number 9, 2018 Mary Ann Liebert, Inc. doi: 10.1089/ars.2017.7118

Noise & Health. (Thomas H. Fay, Ed.). The New York Academy of Medicine, New York, New York. 1991.

Öhrström, E. (2004). Longitudinal surveys on effects of changes in road traffic noise—annoyance, activity disturbances, and psycho-social well-being. *The Journal of the Acoustical Society of America* 115, 719; doi: 10.1121/1.1639333

Roberts, B., and Neitzel, R.L. (2019). Noise exposure limit for children in recreational settings: Review of available evidence. *The Journal of the Acoustical Society of America* 146, 3922; doi: 10.1121/1.5132540.

Rocha, S., Smith, M.G., Witte, M., and Basner, M. (2019). Survey Results of a Pilot Sleep Study Near Atlanta International Airport. *International Journal of Environmental Research and Public Health*, 16, 4321; doi:10.3390/ijerph16224321

Shield, B., and Dockrell, J.E. (2004). External and internal noise surveys of London primary schools. *The Journal of the Acoustical Society of America* 115, 730; doi: 10.1121/1.1635831

Shield, B.M., and Dockrell, J.E. (2008). The effects of environmental and classroom noise on the academic attainments of primary school children Citation: *The Journal of the Acoustical Society of America* 123, 133; doi: 10.1121/1.2812596

Sloan, R.P. Cardiovascular Effects of Noise, in *Noise & Health*. (Thomas H. Fay, Ed.). The New York Academy of Medicine, New York, New York. 1991. (15-26).

Raymond, L.W. “Neuroendocrine, Immunologic, and Gastrointestinal Effects of Noise, in *Noise & Health*. (Thomas H. Fay, Ed.). The New York Academy of Medicine, New York, New York. 1991. (27-40).

Pollak, C.P. “The Effect of Noise On Sleep”, in *Noise & Health*. (Thomas H. Fay, Ed.). The New York Academy of Medicine, New York, New York. 1991. (41-60).

van Gerven, P.W.M., Voss, H., van Boxtel, M.P.J., Janssen, S.A., and Miedema, H.M.E. (2009). Annoyance from environmental noise across the lifespan. *Journal of the Acoustical Society of America*. 126, 1, 187-194.

van Kamp, I., Simon, S., Notley, H., Baliatsas, C., and van Kempen, E. (2020). Evidence Relating to Environmental Noise Exposure and Annoyance, Sleep Disturbance, Cardio-Vascular

and Metabolic Health Outcomes in the Context of IGCB (N): A Scoping Review of New Evidence. *International Journal of Environmental Research and Public Health*

Walker, E.D., Brammer, A., Cherniack, M.G., Laden, F., and Cavallari, J.M. (2016). Cardiovascular and Stress Responses to Short-Term Noise Exposures—A Panel Study in Healthy Males, *Environmental Research*. 150: 391–397.

Wyon, D. (1968). Studies of children under imposed noise and heat stress. *Ergonomics*, 13: 598-612.

WHO Reports:

World Health Organization. *Environmental Noise Guidelines for the European Region*; WHO: Geneva, Switzerland, 2018.

Basner, M.; McGuire, S. WHO Environmental Noise Guidelines for the European Region: A Systematic Review on Environmental Noise and Effects on Sleep. *International Journal of Environmental Research and Public Health*, 2018, 15, 519.

Guski, R.; Schreckenberg, D.; Schuemer, R. WHO Environmental Noise Guidelines for the European Region: A Systematic Review on Environmental Noise and Annoyance. *International Journal of Environmental Research and Public Health*, 2017, 14, 1539.

Clark, C.; Paunovic, K. WHO Environmental Noise Guidelines for the European Region: A Systematic Review on Environmental Noise and Cognition. *International Journal of Environmental Research and Public Health*, 2018, 15, 285.

Smith, M.G., Croy, I., Ogren, M., Hammar, O., Lindberg, E., and Persson Waye, K. (2017). Physiological effects of railway vibration and noise on sleep, *Journal of the Acoustical Association of America*, 141 (5). 3262-3269.

Van Kempen, E.E.M.M.; Casas, M.; Pershagen, G.; Foraster, M. *Cardiovascular and Metabolic Effects of Environmental Noise. Systematic Evidence Review in the Framework of the Development of the WHO Environmental Noise Guidelines for the European Region*; RIVM: Bilthoven, The Netherlands, 2017.

Van Kempen, E.; Casas, M.; Pershagen, G.; Foraster, M. (2018). WHO Environmental Noise Guidelines for the European Region: A Systematic Review on Environmental Noise and Cardiovascular and Metabolic Effects: A Summary. *International Journal of Environmental Research and Public Health*, 15, 379.

Nieuwenhuijsen, M.; Ristovska, G.; Dadvand, P. WHO Environmental Noise Guidelines for the European Region: A Systematic Review on Environmental Noise and Adverse Birth Outcomes. *International Journal of Environmental Research and Public Health*, 2017, 14, 1252.

Sliwinska-Kowalska, M.; Zaborowski, K. WHO Environmental Noise Guidelines for the European Region: A Systematic Review on Environmental Noise and Permanent Hearing Loss and Tinnitus. *International Journal of Environmental Research and Public Health*, 2017, 14, 1139.